

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
18 December 2003 (18.12.2003)

(10) International Publication Number  
**PCT**  
**WO 03/104182 A1**

- (51) International Patent Classification<sup>7</sup>: **C07C 69/54**, 67/03, G03F 7/039
- (74) Agent: **JANG, Seongku**; 19th Fl., KEC Building, #275-7, Yangjae-dong, Seocho-ku, Seoul 137-130 (KR).
- (21) International Application Number: **PCT/KR03/01151**
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 11 June 2003 (11.06.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
10-2002-0032554 11 June 2002 (11.06.2002) KR
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SI, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **ENF TECHNOLOGY CO., LTD.** [KR/KR]; #205-5, Non-hyun-dong, Kangnam-gu, Seoul 135-010 (KR).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **BAE, Eun Hyoung** [KR/KR]; #721-23, Yaeum-3-dong, Nam-gu, Ulsan 680-043 (KR). **SONG, Young Bae** [KR/KR]; #304-2, Yeompo-dong, Buk-gu, Ulsan 683-040 (KR). **OH, Dong Ju** [KR/KR]; #134-3, Doma-1-dong, Seo-gu, Daejeon 302-161 (KR).
- Published:  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: 2-ALKOXYALKYL-2-ADAMANTYL (METH)ACRYLATE AND METHOD FOR PREPARING SAME



(57) Abstract: 2-Alkoxyalkyl-2-adamantyl (meth)acrylate and a method for preparing same are provided.

## 2-ALKOXYALKYL-2-ADAMANTYL (METH)ACRYLATE AND METHOD FOR PREPARING SAME

### Field of the Invention

5

The present invention relates to a novel adamantane derivative, 2-alkoxyalkyl-2-adamantyl (meth)acrylate, and a method for preparing same.

### Background of the Invention

10

Adamantane derivatives having a vinyl substituent have been used in the production of photocurable resins having excellent optical and mechanical properties suitable for applications in such fields as adhesive, printing ink, photosensitive resin, and optical fiber coat. For example, Japanese Laid-open Patent Publication No. Hei9-73173 discloses 2-adamantyl (meth)acrylate useful for the production of chemical amplification resist; and European Laid-open Patent Publication No. 1 000 924 A1, an acid-sensitive polymer having adamantyl moieties which has sufficiently high etching resistance for providing a finer line pattern.

20

Adamantyl derivatives have also been used as drug intermediates. For example, US. patent No. 6,127,415 discloses apoptosis-inducing adamantyl derivatives and their usage as antitumor agents; and Korean Laid-open Patent Publication No. 1997-42511, a pharmaceutical composition comprising an adamantyl derivative which is active against dermatological diseases.

25

### Summary of the Invention

Accordingly, it is an object of the present invention to provide a novel adamantane derivative which is useful as a monomer for the production of a photocurable resin or as a drug intermediate.

30

It is another object of the present invention to provide a method for preparing the adamantane derivative.

### **Brief Description of the Drawings**

5

The above and other objects and features of the present invention will become apparent from the following description of the invention taken in conjunction with the following accompanying drawings, which respectively show:


10 FIG. 1 : an FT-IR spectrum of 2-(4-methoxybutyl)-2-adamantanol produced in Example 1 of the present invention;

FIG. 2 : an NMR spectrum of 2-(4-methoxybutyl)-2-adamantyl acrylate produced in Example 1 of the present invention; and

15 FIG. 3 : an NMR spectrum of 2-(4-methoxybutyl)-2-adamantyl methacrylate produced in Example 2 of the present invention.

### **Detailed Description of the Invention**

20 In accordance with one aspect of the present invention, there is provided 2-alkoxyalkyl-2-adamantyl (meth)acrylate represented by the following formula I:



wherein, R<sup>1</sup> is hydrogen, C<sub>1-4</sub> alkyl or C<sub>3-8</sub> cycloalkyl; R<sup>2</sup> is hydrogen or methyl;  
30 and n is an integer.

Preferably,  $R^1$  is methyl and  $n$  is an integer in the range of 1 to 4.

The inventive 2-adamantyl (meth)acrylate is useful as a resist material suitable for short-wave exposure source (ArF excimer laser) since it is an acid-sensitive compound containing an alkali-soluble group protected with an alicyclic hydrocarbon group.

In accordance with another aspect of the present, there is provided a method of preparing the compound of formula I comprising:

a) reacting 2-adamantanone with a Grignard reagent of formula  $R^1O(CH_2)_nMgX$  to obtain a 2-alkoxyalkyl-2-adamantanol, the alkoxyalkyl being  $R^1O(CH_2)_n$ -; and

b) reacting the 2-alkoxyalkyl-2-adamantanol with (meth)acryloyl chloride to obtain 2-alkoxyalkyl-2-adamantyl (meth)acrylate, wherein  $X$  is chlorine or bromine.


Preferably, step b) is performed in the presence of a basic reagent.

In accordance with a preferred embodiment of the present invention, the method of preparing the compound may further comprise a step of reacting alkoxyalkyl halide with magnesium to obtain alkoxyalkyl Grignard reagent before step a).

In the preparation method of the compound according to the present invention, a separation process may be performed after each step is completed. However, the separation process may be performed after all steps are performed in situ. Accordingly, synthesized Grignard reagent may be employed in step a) without further separation.


A process for producing 2-alkoxyalkyl-2-adamantyl (meth)acrylate according to the present invention will now be described in detail.

As shown in reaction scheme (1), an alkoxyalkyl halide is reacted with magnesium to synthesize an alkoxyalkyl magnesium halide (an alkoxyalkyl reagent):




5 wherein,  $\text{R}^1$  is hydrogen,  $\text{C}_{1-4}$  alkyl or  $\text{C}_{3-8}$  cycloalkyl; X is Cl or Br; and n is an integer.

Next, as shown in reaction scheme (2), adamantanone and the alkoxyalkyl Grignard reagent are reacted, to synthesize a 2-alkoxyalkyl-2-adamantanol:



15 wherein,  $\text{R}^1$  and n are the same as defined above.

Finally, as shown in reaction scheme (3), the 2-alkoxyalkyl-2-adamantanol is reacted with acryloyl or methacryloyl chloride to obtain 2-alkoxyalkyl-2-adamantyl (meth)acrylate:



wherein,  $\text{R}^1$ ,  $\text{R}^2$  and n are the same as defined above.

The alkoxyalkyl Grignard reagent is employed preferably in an amount ranging from 1.2 to 2.0 equivalents based on the amount of 2-adamantanone, and  
 30 (meth)acryloyl chloride is employed preferably in an amount ranging from 1.2 to

2.0 mole per mole of 2-alkoxyalkyl-2-adamantanol.

Preferably, the basic reagent that may be used in the present invention is a tertiary amine such as triethyl amine or pyridine and is employed in an amount ranging from 1.4 to 2.2 mole per mole of 2-alkoxyalkyl-2-adamantanol.

5 In accordance with a preferred embodiment of the present invention, the reaction for synthesizing 2-alkoxyalkyl-2-adamantanol is performed at a temperature ranging from  $-20$  to  $0^{\circ}\text{C}$ , for a period ranging from 2 to 3 hours; and the reaction for synthesizing 2-alkoxyalkyl-2-adamantyl (meth)acrylate, at a temperature ranging from  $-20$  to  $0^{\circ}\text{C}$ , for a period ranging from 2 to 15 hours.

10 The present invention is described in more detail below by referring to the following Examples, and the Examples are given for the purpose of illustration only, and are not intended to limit the scope of the invention.

Example 1: Preparation of 2-(4-methoxybutyl)-2-adamantyl acrylate

15

A. Synthesis of 4-methoxybutyl magnesium chloride (4-methoxybutyl Grignard reagent)

52 g of 4-methoxy-1-butanol (0.5 mol) and 40 g of pyridine (0.5 mol) were placed into a 1 liter three-necked flask and then cooled to  $0^{\circ}\text{C}$ . Thionyl chloride (119 g, 1.0 mol) was added slowly thereto using a dropping funnel. 20 The mixture was refluxed for about 1 hour, after poured into water to destroy completion of the reaction, excess thionyl chloride extracted with 300 ml of methylene chloride. Then, the organic layer was washed with 5% sodium hydroxide and dried over magnesium sulfate to obtain 4-methoxybutyl chloride 25 (yield: 80%).

26.7 g of magnesium turnings (1.1 mol) was dispersed in 500ml of anhydrous THF, and then, a small amount of iodine and a drop of 4-methoxybutyl chloride were added thereto, followed by heating up to  $40^{\circ}\text{C}$  to activate magnesium. After the color of the solution changed from brown to 30 colorless, the mixture was cooled and then 108.57 g of the 4-methoxybutyl

chloride (1.0 mol) was added slowly thereto. After completion of the addition, the reaction mixture was warmed, and refluxed for about 1 hour to obtain a 4-methoxybutylmagnesium chloride solution.

5           B. Synthesis of 2-(4-methoxybutyl)-2-adamantanol

The 4-methoxybutylmagnesium chloride solution obtained in step A in an amount corresponding to 1.0 mole of 4-methoxybutyl chloride was placed in a 1 liter flask using a cannular and cooled to 0 °C. 120.17 g of 2-adamantanone (0.8 mol) was added slowly thereto using a dropping funnel, and stirred at room  
10    temperature for about 12 hours. After completion of the reaction, excess THF was removed using a rotary evaporator, the resulting residue was poured into water, the solution was neutralized with dilute sulfuric acid, extracted using diethyl ether, and dried over magnesium sulfate. The crude product was re-crystallized from n-hexane-methylene chloride to obtain 2-(4-methoxybutyl)-2-  
15    adamantanol (yield: 80 %).

FT-IR (KBr;  $\text{cm}^{-1}$ ): 3431, 2902, 1449.5, 1112.5, 941.5, 611.

C. Synthesis of 2-(4-methoxybutyl)-2-adamantyl acrylate

44.87 g of the 2-(4-methoxybutyl)-2-adamantanol (0.2 mol) obtained in  
20    step B and 32.38 g of triethylamine (0.32 mol) were dissolved in 300 ml of THF and then 25.34 g of acryloyl chloride (0.28 mol) was added slowly thereto using a dropping funnel. Then, the reaction was stirred at room temperature for about 12 hours.

After completion of the reaction, excess THF was removed using a  
25    rotary evaporator, and the resulting mixture was poured into water. The solution was neutralized with dilute sulfuric acid, extracted using diethyl ether, and the ether layer was dried over magnesium sulfate. After removing diethyl ether, the crude product was vacuum distilled to obtain 2-(4-methoxybutyl)-2-adamantyl acrylate (yield: 80%).

30            $^1\text{H}$ -NMR ( $\text{CDCl}_3$ ; ppm): 6.25(1H, d), 6.1(1H, dd), 5.9(1H, d), 3.3(2H, t),

3.2(3H, S), 2.3-1.2(m).

Example 2: Preparation of 2-(4-methoxybutyl)-2-adamantyl methacrylate

5           The title compound was prepared in the same manner as in Example 1, except that 29.27 g of methacryloyl chloride (0.28 mol) was used instead of 25.34 g of acryloyl chloride (yield: 85%).

          H-NMR (CDCl<sub>3</sub>; ppm): 6.1(1H, d), 5.5(1H, d), 3.3(2H, t), 3.2(3H, S), 2.3-1.2(m).

10

          2-Alkoxyalkyl-2-adamantyl (meth)acrylate according to the present invention can be used for various applications e.g. as a monomer of a photocurable resin, a drug intermediate and the like. Also, the inventive method for preparing 2-alkoxyalkyl-2-adamantyl (meth)acrylate is simple, a high yield  
15   process suitable for commercial-scale production.


          While the invention has been described with respect to the above specific embodiments, it should be recognized that various modifications and changes may be made to the invention by those skilled in the art which also fall within the scope of the invention as defined by the appended claims.

20



**What is claimed is :**

1. A compound of formula I:



wherein,  $R^1$  is hydrogen,  $C_{1-4}$  alkyl or  $C_{3-8}$  cycloalkyl;  $R^2$  is hydrogen or methyl; and  $n$  is an integer.

- 15 2. The compound of claim 1, wherein  $n$  is an integer in the range of 1 to 4.

3. A method of preparing the compound of claim 1, comprising:

- a) reacting 2-adamantanone with a Grignard reagent of formula  $R^1O(CH_2)_nMgX$  to obtain a 2-alkoxyalkyl-2-adamantanol, the alkoxyalkyl being  $R^1O(CH_2)_n-$ ; and
- 20

- b) reacting the 2-alkoxyalkyl-2-adamantanol with (meth)acryloyl chloride to obtain 2-alkoxyalkyl-2-adamantyl (meth)acrylate, wherein  $X$  is chlorine or bromine.


25

4. The method of claim 3, wherein step b) is performed in the presence of a basic reagent.

5. The method of claim 3, wherein in step a) 2-alkoxyalkyl-2-adamantanol is 2-(4-methoxybutyl)-2-adamantanol.
- 30


1/3

FIG. 1



2/3


FIG. 2



SUBSTITUTE SHEET (RULE 26)

3/3

FIG. 3



SUBSTITUTE SHEET (RULE 26)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR03/01151

**A. CLASSIFICATION OF SUBJECT MATTER****IPC7 C07C 69/54, 67/03, G03F 7/039**

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

C07C, G03F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
Korean Patents and Applications for inventions since 1975Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
STN(REG,CA)**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

| Category* | Citation of document, with indication, where appropriate, of the relevant passages                      | Relevant to claim No. |
|-----------|---|-----------------------|
| X         | KR 2001- 81855 A(Chem Search Corp.) 29 Aug. 2001<br>see claim 1-3                                       | 1-5                   |
| X         | US 6222061 B1(Chem Search Corp.) 24 Apr. 2001<br>see column2 line45- column3 line10                     | 1-5                   |
| X         | EP 1127870 A1(Shipley Co. Llc) 29 Aug. 2001<br>see page3 line35-page4 line 23, page6 Reaction scheme1-5 | 1-5                   |
| X         | EP 1020767 A1(Sumitomo Chemical Co. Ltd.)19 Jul. 2000<br>see example2                                   | 1-5                   |
| X         | US 6013416 A(Fujitsu Ltd.) 11 Jan. 2000<br>see column27 chemical formula( X X X )                       | 1-2                   |

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 JULY 2003 (23.07.2003)

Date of mailing of the international search report

24 JULY 2003 (24.07.2003)

Name and mailing address of the ISA/KR



Korean Intellectual Property Office  
920 Dunsan-dong, Seo-gu, Daejeon 302-701,  
Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

KANG, Jeon Kwan

Telephone No. 82-42-481-5553



Form PCT/ISA/210 (second sheet) (July 1998)

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR03/01151

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s)                         | Publication<br>date                    |
|---|---------------------|--|--|
| KR 2001- 81855 A                          | 29-08-2001          | NONE   |  |
| US 6222061 B1                             | 24-04-2001          | KR 2001-81853 A<br>JP 13-233834 A<br>EP 1125917 A1 | 29-08-2001<br>28-08-2001<br>22-08-2001 |
| EP 1127870 A1                             | 29-08-2001          | KR 2001-85618 A<br>JP 14-3447 A2                   | 07-09-2001<br>09-01-2002               |
| EP 1020767 A1                             | 19-07-2000          | JP 12-275845 A2<br>US 6383713 B1<br>CN 1261171 A   | 06-10-2000<br>07-05-2002<br>26-07-2000 |
| US 6013416 A                              | 11-01-2000          | KR 206664 B1<br>JP 9-73173 A<br>DE 19626003 A1     | 01-07-1999<br>18-03-1997<br>02-01-1997 |
| -----                                     | End of line         | -----  | -----                                  |